

A Binary Origin for Blue Stragglers in Globular Clusters

Christian Knigge¹, Nathan Leigh², & Alison Sills²

¹*University of Southampton, School of Physics and Astronomy, Southampton SO17 1BJ, UK*

²*McMaster University, Department of Physics and Astronomy, 1280 Main Street West, Hamilton, ON, L8S 4M1, Canada*

Blue stragglers in globular clusters are abnormally massive stars that should have evolved off the stellar main sequence long ago. There are two known processes that can create these objects: direct stellar collisions¹ or binary evolution². However, the relative importances of these processes have remained unclear. In particular, the total number of blue stragglers found in a given cluster does not seem to correlate with the predicted collision rate^{3,4}, providing indirect support for the binary scenario. Yet the radial distributions of blue stragglers in many clusters are bimodal, with a dominant central peak^{5–7}. This has been interpreted as an indication that collisions do dominate blue straggler production, at least in the high density cluster cores^{7,8}. Here, we show that there is a clear, but sub-linear, correlation between the number of blue stragglers found in a cluster core and the total stellar mass contained within it. This is the strongest and most direct evidence to date that most blue stragglers, even those found in cluster cores, are the progeny of binary systems. However, we also point out that the parent binaries may themselves have been affected by dynamical encounters. This may be the key to reconciling all of the seemingly conflicting results found to date.

In both blue straggler (BS) formation scenarios, the numbers and properties of the resultant BSs can shed light on the dynamical history of their host clusters. The conclusions drawn from the observed BS populations depend on which formation mechanism is dominant. Unfortunately, even though a number of potential formation tracers have been proposed, a simple observational distinction between collisional and binary BSs has yet to be found^{9–14}. In general, it is impossible to ascertain whether an individual BS was formed via a stellar collision or a binary merger.

However, it should be possible to determine the dominant formation channel statistically, by studying the relationship between the size of a BS population and the properties of its host cluster. Here, we focus on BSs found in the cores of clusters, where collisions should be most frequent. If most BSs are formed as a result of single-single collisions, their number in a particular cluster core should be approximately $N_{BSS, coll} \simeq \tau_{BSS}/\tau_{coll}$, where τ_{BSS} is the typical BS lifetime, and τ_{coll} is the time-scale between single-single stellar collisions in the core. Conversely, if most core BSs are the progeny of binary stars, their number may

be expected to scale simply as $N_{BSS,bin} \propto f_{bin} M_{core}$, where f_{bin} is the binary fraction in the core and M_{core} is the total stellar mass contained in the core. These predictions are for the simplest versions of the competing scenarios, but note that more complex, intermediate channels could also contribute (e.g. BSs produced as a result of dynamical encounters involving binary stars). We will return to this subtlety below.

In order to test these simple predictions, we used our existing data base of core BS counts⁴, which are derived from a large set of *HST*-based colour-magnitude diagrams¹⁵. Figure 1 confirms that there is no global correlation between the observed core BS numbers and those predicted by the collision scenario^{3,4}. In particular, the predicted numbers are very different for high- and low-density clusters, which is in conflict with the observations. By contrast, Figure 2 shows that the observed numbers do correlate strongly with the cluster core masses, as expected if most BSs are the progeny of binary stars. A power law fit suggests $N_{BSS} \propto M^{0.38 \pm 0.04}$.

When consideration is restricted to dense cores ($\rho_0 \gtrsim 10^4 \text{ M}_\odot \text{ pc}^{-3}$), the collisional predictions are, in fact, significantly correlated with observed BS numbers (see the black points in Figure 1). However, even for these dense cores, the correlation with core mass remains stronger (Figure 2). Thus not only is core mass the only successful global predictor for core BS numbers, but it remains a better predictor than collision rate even for dense clusters.

One way to distinguish even more clearly between collisional and binary scenarios is to isolate the key cluster parameters that drive the observed correlations and quantify their impact on the observed sizes of the core BS populations. For example, even though collision rate and core mass are correlated parameters, only collision rate depends fundamentally on a cluster's central velocity dispersion. We therefore carried out a direct fit to the observed core BS numbers, using the generalized model $N_{BSS} = K r_c^\alpha \rho_0^\beta \sigma_0^\gamma$, where K , α , β and γ are all treated as free parameters. For purely collisional BS formation, we expect $\alpha \simeq 3$, $\beta \simeq 2$ and $\gamma \simeq -1$, whereas pure binary formation implies $\alpha \simeq 3$, $\beta \simeq 1$ and $\gamma \simeq 0$. These relationships assume constant binary fraction across clusters; we will return to this assumption below. Allowing for a more general power-law dependence on collision rate or core mass, we predict $\alpha \simeq 1.5\beta \simeq -3\gamma$ for collisional BS formation, and $\alpha \simeq 3\beta$ with $\gamma \simeq 0$ for the binary scenario.

Figure 3 shows the results of this fit. The best-fit power law indices are $\alpha = 1.47 \pm 0.29$, $\beta = 0.55 \pm 0.15$ and $\gamma = -0.40 \pm 0.27$. These coefficients are consistent with a power-law dependence on M_{core} , but not with one on collision rate. (Note that since r_c , ρ_0 and σ are correlated parameters, α , β and γ are also correlated.) There is no evidence for a significant dependence of BS numbers on velocity dispersion. A fit to the dense clusters only yielded results which were consistent with the values obtained for the complete cluster sample.

We find that BS numbers depend strongly on core mass, but not on collision rate. We argue that this is strong evidence for a binary, as opposed to a collision, origin for blue stragglers in globular clusters. We note that the core mass dependence can be approximated as $N_{BSS} \propto M_{core}^{\delta}$ with $\delta \simeq 0.4 - 0.5$. The sub-linear dependence of BS numbers on core mass is interesting. For ordinary populations of single stars, one would expect a roughly linear dependence, and we have verified that this is the case for the red giant and horizontal branch populations in our cluster sample. In the context of the binary scenario for BSs, the simplest way to account for this sub-linearity is to posit that core binary fractions decrease with increasing core mass, i.e. $f_{bin} \propto M_{core}^{-\epsilon}$ with $\epsilon \simeq 0.5 - 0.6$. This is a testable prediction. There have been two recent efforts to estimate core binary fractions in a consistent way from homogeneous, deep HST/ACS observations of globular clusters^{25,26}. One of these²⁵ is restricted to a set of 13 low-density clusters (Sample 1). The other²⁶ is still preliminary, but based on a larger set of 35 clusters that span a wider range in density and other dynamical parameters (Sample 2). Promisingly, from our perspective, a correlation between BS frequency and core binary fraction has already been reported for Sample 1²⁷, and an anti-correlation between core binary fraction and total cluster mass has been reported for Sample 2²⁶. Since Sample 1 spans only a narrow dynamic range, and the binary fractions reported for Sample 2 are only available in graphical form, we could only make a preliminary attempt to infer the dependence of f_{bin} on M_{core} from these samples. Nevertheless, our power law estimate is $f_{bin} \propto M_{core}^{-0.35}$ for both samples, close to the predicted relation, albeit with considerable scatter. Thus the number of BSs found in the cores of globular clusters scales roughly as $N_{BSS} \propto f_{bin} M_{core}$, just as expected if most core BSs are descended from binary systems.

There are several ways to build on the analysis presented above. First, binary-single and binary-binary interactions may actually produce more blue stragglers in clusters than single-single collisions because the interaction cross sections are larger. The expected number of binary-single [binary-binary] collisions scales like the number of single-single collisions times f_b [f_b^2]. A better determination of the dependence of binary fractions on cluster parameters is clearly warranted to test these predictions fully. However, based on the preliminary analysis of the observed binary fractions described above, it currently seems unlikely that binary-single and binary-binary interaction numbers can produce the observed scaling of blue straggler numbers with cluster parameters.

Second, the effects of mass segregation have been ignored in our analytical predictions. Massive stars at any particular location in a cluster will sink to the core on roughly the local relaxation time. Thus some of the BSs found today in cluster cores may actually have been formed outside the core^{7,8,28}. Moreover, clusters evolve, so the present-day cluster parameters may not always be close approximations to what should ideally be suitably-weighted averages over the BS lifetime. As a first test, we have split our cluster sample in objects with half-mass relaxation times smaller and greater than the blue straggler lifetime. If mass segregation were responsible for masking an underlying trend with collision rate,

we may expect the long relaxation timescale clusters to reveal this trend, while the short relaxation time clusters should exhibit enhanced BS numbers (at a given collision rate). We find no evidence for either of these effects. It is also worth noting that our result for core BSs is actually consistent with the behaviour of the total (core+peripheral) BS populations, which also seem to show a sub-linear correlation with total cluster mass²⁸.

Our results do not imply that dynamical interactions are unimportant to BS formation. The properties of the core binary populations from which BSs are probably descended may themselves be strongly affected by dynamical encounters. There is a wide spectrum of possible paths within the family of binary channels. While some 'binary BSs' may be the direct descendants of primordial binaries that evolved essentially in isolation, others may exist only as a result of multiple strong interactions and exchanges^{29,30}. We are not the first to point out this subtlety (though it has sometimes been obscured by the terminology used). For example, Mapelli et al. note that the collisional BS population invoked in their model could be produced via encounters involving binaries^{7,8}, and the primordial binary channel described by Davies et al. explicitly allows for dynamical interactions²⁸. Our results do, however, argue against the simple scaling of blue straggler numbers with any collision rate. Clearly, a key challenge for future work will be to determine if dynamically active binaries can produce BS numbers consistent with the trends found here.

1. Hills, J. G. & Day, C. A. Stellar Collisions in Globular Clusters. *Astrophys. Lett.* **17**, 87–93 (1976).
2. McCrea, W. H. Extended main-sequence of some stellar clusters. *Mon. Not. R. Astron. Soc.* **128**, 147–155 (1964).
3. Piotto, G. *et al.* Relative Frequencies of Blue Stragglers in Galactic Globular Clusters: Constraints for the Formation Mechanisms. *Astrophys. J. Lett.* **604**, L109–L112 (2004).
4. Leigh, N., Sills, A. & Knigge, C. Where the Blue Stragglers Roam: Searching for a Link between Formation and Environment. *Astrophys. J.* **661**, 210–221 (2007); Erratum: *Astrophys. J.*, **678**, 564–566 (2008).
5. Ferraro, F. R. *et al.* Blue Stragglers in the Galactic globular clusters M3: Evidence for two populations. *Astronom. J.* **106**, 2324–2334 (1993).
6. Ferraro, F. R. *et al.* Discovery of Another Peculiar Radial Distribution of Blue Stragglers in Globular Clusters: The Case of 47 Tucanae. *Astrophys. J.* **603**, 127–134 (2004).
7. Mapelli, M. *et al.* The radial distribution of blue straggler stars and the nature of their progenitors. *Mon. Not. R. Astron. Soc.* **373**, 361–368 (2006).
8. Mapelli, M. *et al.* The Contribution of Primordial Binaries to the Blue Straggler Population in 47 Tucanae. *Astrophys. J. Lett.* **605**, L29–L32 (2004).

9. Benz, W. & Hills, J. G. Three-dimensional hydrodynamical simulations of stellar collisions. I - Equal-mass main-sequence stars. *Astrophys. J.* **323**, 614–628 (1987).
10. Leonard, P. J. T. & Livio, M. The Rotational Rates of Blue Stragglers Produced by Physical Stellar Collisions. *Astrophys. J. Lett.* **447**, L121–L123 (1995).
11. Procter Sills, A., Bailyn, C. D. & Demarque, P. Are Blue Stragglers Mixed during Collisions? *Astrophys. J. Lett.* **455**, L163–L165 (1995).
12. Sandquist, E. L., Bolte, M. & Hernquist, L. Composition Mixing during Blue Straggler Formation and Evolution. *Astrophys. J.* **477**, 335–345 (1997).
13. Sills, A. & Bailyn, C. D. The Distribution of Collisionally Induced Blue Stragglers in the Color-Magnitude Diagram. *Astrophys. J.* **513**, 428–441 (1999).
14. Ferraro, F. R. *et al.* Discovery of Carbon/Oxygen-depleted Blue Straggler Stars in 47 Tucanae: The Chemical Signature of a Mass Transfer Formation Process. *Astrophys. J. Lett.* **647**, L53–L56 (2006).
15. Piotto, G. *et al.* HST color-magnitude diagrams of 74 galactic globular clusters in the HST F439W and F555W bands. *Astron. & Astrophys.* **391**, 945–965 (2002).
16. Gehrels, N. Confidence limits for small numbers of events in astrophysical data. *Astrophys. J.* **303**, 336–346 (1986).
17. Leonard, P. J. T. Stellar collisions in globular clusters and the blue straggler problem. *Astronom. J.* **98**, 217–226 (1989).
18. Harris, W. E. A Catalog of Parameters for Globular Clusters in the Milky Way. *Astronom. J.* **112**, 1487–1488 (1996).
19. McLaughlin, D. E. & van der Marel, R. P. Resolved Massive Star Clusters in the Milky Way and Its Satellites: Brightness Profiles and a Catalog of Fundamental Parameters. *Astrophys. J. Supp.* **161**, 304–360 (2005).
20. Pryor, C. & Meylan, G. Velocity Dispersions for Galactic Globular Clusters. In Djorgovski, S. G. & Meylan, G. (eds.) *Structure and Dynamics of Globular Clusters*, vol. 50 of *Astronomical Society of the Pacific Conference Series*, 357–371 (1993).
21. Webbink, R. F. Structure parameters of galactic globular clusters. In Goodman, J. & Hut, P. (eds.) *Dynamics of Star Clusters*, vol. 113 of *IAU Symposium*, 541–577 (1985).
22. Sills, A. *et al.* Evolution of Stellar Collision Products in Globular Clusters. I. Head-on Collisions. *Astrophys. J.* **487**, 290–303 (1997).
23. Sills, A., Faber, J. A., Lombardi, J. C., Jr., Rasio, F. A. & Warren, A. R. Evolution of Stellar Collision Products in Globular Clusters. II. Off-Axis Collisions. *Astrophys. J.* **548**, 323–334 (2001).

24. Verbunt, F. W. M. Observational evidence for origin of stellar exotica in globular clusters. *Highlights of Astronomy* **14**, 440–441 (2007).
25. Sollima, A., Beccari, G., Ferraro, F. R., Fusi Pecci, F. & Sarajedini, A. The fraction of binary systems in the core of 13 low-density Galactic globular clusters. *Mon. Not. R. Astron. Soc.* **380**, 781–791 (2007).
26. Milone, A. P., Piotto, G., Bedin, L. R. & Sarajedini, A. Photometric binaries in fifty Globular Clusters. In *XXI Century Challenges for Stellar Evolution*, vol. 79 of *Memorie della Società Astronomica Italiana*, in press (2008). [arXiv:0801.3177](#).
27. Sollima, A., Lanzoni, B., Beccari, G., Ferraro, F. R. & Fusi Pecci, F. The correlation between blue straggler and binary fractions in the core of Galactic globular clusters. *Astron. & Astrophys.* **481**, 701–704 (2008).
28. Davies, M. B., Piotto, G. & de Angeli, F. Blue straggler production in globular clusters. *Mon. Not. R. Astron. Soc.* **349**, 129–134 (2004).
29. Knigge, C. *et al.* A Blue Straggler Binary with Three Progenitors in the Core of a Globular Cluster? *Astrophys. J.* **641**, 281–287 (2006).
30. Knigge, C. *et al.* Stellar Exotica in 47 Tucanae. *Astrophys. J.* **in press** (2008). [arXiv:astro-ph/0805.0140](#).

Acknowledgements Research support for C.K. was provided by the UK’s Science and Technology Facilities Council. A.S. and N.L. are supported by the Natural Sciences and Engineering Research Council of Canada. C.K. would like to thank Tom Maccarone for useful discussions.

Author Information Reprints and permissions information is available at npg.nature.com/reprintsandpermissions. The authors declare that they have no competing financial interests. Correspondence and requests for materials should be addressed to C. K. (email: christian@astro.soton.ac.uk).




Figure 1: The observed numbers of core blue stragglers versus the numbers expected from single-single collisions. Black points correspond to high-density clusters ($\rho_0 > 10^4 \text{ M}_\odot \text{ pc}^{-3}$), grey points to low-density ones ($\rho_0 < 10^4 \text{ M}_\odot \text{ pc}^{-3}$). The error bars shown are approximate 68% Poisson confidence intervals¹⁶. There is no correlation between observed and predicted numbers across the full sample (Spearman’s rank test yields $\rho_s = -0.03$, $p = 0.6$, $N = 56$). For the subset of dense clusters, a correlation does exist ($\rho_s = 0.52$, $p = 8 \times 10^{-4}$, $N = 34$). In calculating the predicted numbers, we took the time-scale between collisions to be¹⁷ $\tau_{coll} = 1.1 \times 10^{10} \left(\frac{r_c}{1 \text{ pc}}\right)^{-3} \left(\frac{n_0}{10^3 \text{ pc}^{-3}}\right)^{-2} \left(\frac{\sigma_0}{5 \text{ km s}^{-1}}\right) \left(\frac{M_*}{M_\odot}\right)^{-1} \left(\frac{R_*}{R_\odot}\right)^{-1}$ years. Here r_c is the core radius, n_0 is the central number density in stars per cubic parsec, σ_0 is the central velocity dispersion, M_* is the average stellar mass (taken to be $0.5 \text{ M}_\odot$), and R_* is the average stellar radius ($0.5 \text{ R}_\odot$). Core radii and central luminosity densities were taken from the McMaster Globular Cluster Catalogue¹⁸. Core number densities were estimated by adopting a fixed mass-to-light ratio of 2¹⁹. Central velocity dispersions were taken primarily from the compilation of Pryor & Meylan²⁰, but supplemented with data from Webbink²¹. The typical BS lifetime in a cluster was taken to be 1.5 Gyr^{22,23}. Note that for high-concentration clusters, the observational definition of the core radius can be somewhat arbitrary. However, this should not have a significant impact on our analysis, as long as the associated central densities and velocity dispersions remain reasonable averages over the relevant regions.




Figure 2: The observed numbers of core blue stragglers versus the estimated core masses. Black points correspond to high density clusters ($\rho_0 > 10^4 \text{ M}_\odot \text{ pc}^{-3}$), grey points to low-density ones ($\rho_0 < 10^4 \text{ M}_\odot \text{ pc}^{-3}$). The error bars shown are approximate 68% Poisson confidence intervals¹⁶. BS numbers and core masses are strongly correlated ($\rho_s = 0.71$, $p = 4 \times 10^{-10}$, $N = 56$), as expected if BSs are descended from binary systems. The correlation with core mass also holds for the subset of dense clusters ($\rho_s = 0.84$, $p = 3 \times 10^{-10}$, $N = 34$) and in both cases is stronger than that with collision rate. The dashed line is a power law fit to the full set of clusters, obtained with a method appropriate for this type of data set²⁴. The best-fitting power law is $N_{BSS} \propto M^{0.38 \pm 0.04}$.




Figure 3: Results of fitting the generalized model $N_{BSS} \propto K r_c^\alpha \rho_0^\beta \sigma_0^\gamma$ to the observed BS numbers. The top panels show three projections of the 4-dimensional parameter space. The contours in each panel represent 1σ , 2σ and 3σ joint confidence intervals on the two parameters being considered. Black contours correspond to the full cluster sample, grey contours to the dense cluster sample ($\rho_0 > 10^4 \text{ M}_\odot \text{ pc}^{-3}$). The straight lines in the middle and right panels show the parameter relationships expected for a power law dependence on collision rate (dashed lines) and core mass (solid line). The best fit power law indices are $\alpha = 1.47 \pm 0.29$, $\beta = 0.55 \pm 0.15$ and $\gamma = -0.40 \pm 0.27$. The bottom panel shows observed BS numbers vs the best-fit model prediction for the full sample. Black points correspond to high density clusters ($\rho_0 > 10^4 \text{ M}_\odot \text{ pc}^{-3}$), grey points to low-density ones ($\rho_0 < 10^4 \text{ M}_\odot \text{ pc}^{-3}$). The error bars shown are approximate 68% Poisson confidence intervals¹⁶. The dashed line marks the identity relation, i.e. $N_{BSS}(\text{observed}) = N_{BSS}(\text{predicted})$.

We also carried out a separate fit to the subset of dense clusters ($\rho_0 > 10^4 \text{ M}_\odot \text{ pc}^{-3}$). This yielded $\alpha = 1.59 \pm 0.33$, $\beta = 0.63 \pm 0.19$ and $\gamma = -0.28 \pm 0.31$, consistent with the values obtained for the complete cluster sample. Note that, even within this subset, there is no evidence for a dependence of BS numbers on σ_0 .